

Genetic and Perinatal Risk Factors for Asthma Onset and Severity: A Review and Theoretical Analysis

Michael B. Bracken^{1,2}, Kathleen Belanger¹, William O. Cookson³, Elizabeth Triche¹, David C. Christiani⁴, and Brian P. Leaderer¹

¹ Center for Perinatal, Pediatric and Environmental Epidemiology, Department of Epidemiology and Public Health, Yale University School of Medicine, New Haven, CT.

² Department of Obstetrics and Gynecology, Yale University School of Medicine, New Haven, CT.

³ The Wellcome Trust, University of Oxford, Oxford, United Kingdom.

⁴ Occupational Health Program, Harvard School of Public Health, Cambridge, MA.

Received for publication July 24, 2002; accepted for publication January 3, 2003.

Abbreviations: AR, adrenergic receptor; Ig, immunoglobulin; IL, interleukin; TNF, tumor necrosis factor.

INTRODUCTION

Asthma is a major chronic disease, and several studies indicate that it is on the rise worldwide (1). A recent report (2) from the Centers for Disease Control and Prevention estimated that the prevalence of self-reported asthma in the United States rose 75 percent from 1980 to 1994, with 17.3 million asthmatics in 1998 (3). In 2000, asthma accounted for more than 11.2 million medical visits, including 1.8 million to emergency rooms (4, 5). Asthma is characterized by lung inflammation, reversible airflow obstruction, and enhanced airway responsiveness to a variety of environmental stimuli and is a phenotypically heterogeneous disorder with variable disease expression.

Asthma has a considerably greater impact on Hispanics and African Americans than on Whites in the United States (2, 6–12). Compared with Whites, African-American children have higher (1.1–1.7 times) asthma prevalence rates (2, 13–18), 2–3.5 times the hospital admission rate for asthma (2, 19–23), and approximately 2–5 times the asthma mortality rate (2, 22, 24–26). Point prevalence asthma rates of 11.2 percent and cumulative prevalence rates (ever had asthma) of 20.1 percent are reported for Puerto Rican children, the highest for any ethnic group in the United States (12, 27).

Asthma increasingly has been diagnosed in young children, starting in the 1970s (28) and continuing through the last two decades (29–31), but precise rates cannot be readily obtained because of the difficulty in diagnosing asthma in very young children (32). Wheezing is often used as a surro-

gate measure but is unreliable; in a Tucson, Arizona, cohort, by age 3 years, 19.9 percent of the children had at least one lower respiratory tract illness with wheeze but were no longer wheezing at age 6 years, 15.0 percent did not wheeze before age 3 years but did so at age 6 years, and 49.5 percent wheezed by age 6 years (33). Wheeze has been reported in the winter of the first year of life in 33 percent of infants (34) and to occur for 30 or more days in one third of infants who do wheeze (35). The difficulty in diagnosing childhood asthma has led to suggestions that the increase is in milder symptoms only and that some children may be treated inappropriately (29, 36).

Despite a considerable literature on risk factors for asthma onset and severity in children, very little is known about possible intrauterine influences, particularly how these factors interact with the genotype to sensitize the fetus to allergen exposure in infancy. There have been several recent reviews of candidate genes, but they often have excluded consideration of those environmental risk factors likely to play a role in gene-environment interactions.

In this review, we first discuss some of the major candidate genes currently thought to play a role in affecting susceptibility to allergen sensitization, inflammation and tissue damage, and asthma symptoms and bronchial hyperreactivity. We then consider perinatal risk factors, including intrauterine exposure and influence of the fetal environment. We summarize the literature regarding lactation and diet, early neonatal exposure, and environmental risk factors. Finally, we propose a model that describes the possible interplay of these factors in a plausible temporal sequence.

GENETIC FACTORS

Although environmental factors are clearly important determinants of asthma, numerous studies have revealed that asthma has a strong genetic component but does not follow monogenic patterns of inheritance (37–39). For a long time, asthma has been known to cluster in families, and family studies were the first to suggest that the disease was genetically inherited. More recent family studies found, for example, a 60 percent increased risk of atopy when both parents were affected (40), and the odds of asthma in a child increased from 3 when one parent was affected to 6 when both were (41). Maternal asthma appears to be more influential than paternal asthma (41, 42), particularly in children less than age 5 years (41). While family studies point to the likely importance of a genetic etiology, these studies do not definitively delineate genetic from environmental risks because of shared environments in families.

Twin studies were among the earliest to demonstrate the importance of genetic factors in the etiology of asthma. One of these, conducted in Sweden (43), reported concordance rates for self-reported asthma of 19.0 percent in monozygotic and 4.8 percent in dizygotic twins. There have been many replications of this finding. Current twin studies confirm the importance of both genetic and environmental factors by comparing the concordance rates in monozygotic versus dizygotic twins from the same population, who are at increased risk of asthma because of parental atopy (44). In Finnish Twin Studies, 87 percent of the variation in susceptibility to asthma was owed to genetic factors in families with at least one asthmatic parent. Among families in which neither parent was asthmatic, the development of asthma was explained entirely by environmental risk factors (45, 46). Twin studies permit analysis of environmental risk factors, independent of genetic factors, without necessarily knowing the specific genes involved (47–50).

The strategy for identifying candidate genes offers opportunities to further specify persons at increased risk for susceptibility to allergic sensitization (atopy), inflammation, bronchial hyperreactivity, and severity of asthma symptoms. Confirming the importance of candidate asthma genes will enable development of new diagnostic and therapeutic tools and of prevention and allergen avoidance strategies. Identification of candidate genes for asthma will also permit more precise elucidation of environmental risk factors operating at different stages of asthma development. Nonetheless, careful analysis and interpretation of these studies is required. Three explanations are possible for an association between a candidate gene and disease (51): 1) the candidate allele is the relevant mutation in the disease gene; 2) the allele is positioned very close to the disease gene (linkage disequilibrium); or 3) the association is due to confounding by the allele frequency being higher in population subgroups in which disease frequency is also higher (population admixture). Diseases with a complex genetic origin, such as asthma, also may be characterized by pleiotropy (the same genotype has different phenotypes), genetic heterogeneity (the same phenotype results from different polymorphisms), and incomplete penetrance (the same polymorphism does not always produce the same phenotype).

Multiple regions of the human genome likely to contain susceptibility genes for asthma and associated phenotypes have been reported from candidate-gene approaches and genome-wide screening studies (52–54). For a candidate gene to potentially be important in the disease, a number of criteria must be met. First, the gene protein product must be relevant to the pathophysiology of the disease. Second, the gene must contain mutations within either the coding region or the regulatory regions controlling gene expression; these mutations need to be functionally relevant. Demonstration of functional relevance for a mutation is particularly important given the high rate of polymorphic variation within the human genome, estimated to be about 1 in 1,000 base pairs in coding DNA and about 1 in 500 base pairs in noncoding DNA. Third, functionally relevant mutations should demonstrate association and/or linkage with an appropriate phenotype. Finally, for a mutation to contribute to disease risk in a population, it must be relatively common: rare mutations may greatly increase the risk of developing asthma in individual families but are unlikely to be important in determining the population risk as a whole. However, it follows that the effects of common polymorphisms may be relatively small; if major deleterious consequences occurred in persons with a given polymorphism, it would soon be lost from the population.

Genes for allergic sensitization

Interleukin (IL)-4. Genetic variants in the promoter region of the *IL-4* gene (55) have been related to elevated immunoglobulin (Ig)E levels. The polymorphism at –589 involves a C→T substitution in the promoter region on chromosome 5q31, resulting in increased responsiveness to *IL-4* (e.g., by enhanced IgE production). This locus has been associated with asthma diagnosis in some studies (56, 57). In an Australian population ($n = 1,004$), Walley and Cookson (57) reported positive associations between the *IL-4* promoter polymorphism and specific IgE to dust mite and clinical symptoms of wheeze but could not duplicate these results in a smaller ($n = 183$) English population.

IL-13. Polymorphisms within the *IL-13* gene are associated with high IgE levels and with the presence of asthma (58). *IL4-R α* on chromosome 16 is a shared component of the receptor for both *IL-4* and *IL-13*, and polymorphisms in this gene are also associated with asthma and atopy (59). It is of interest that different asthma-associated traits are associated with individual polymorphisms that affect splicing of *IL4-R α* (60, 61), illustrating the complexity of mechanisms that may vary the actions of a single gene. Gene-gene interactions rarely have been studied, but recently an interaction between polymorphisms in *IL4-R α* and *IL-13* was reported to increase the risk of asthma fivefold (62).

Innate immunity is becoming recognized as equally as important as specific immunity in the response to mucosal and skin injury. The innate immune system contains many molecules that recognize signals of infection, such as components of the bacterial wall and methylated bacterial DNA. These molecules also up-regulate the specific immune system and may enhance IgE responses. *CD14* is a receptor for bacterial lipopolysaccharide. Polymorphism in the *CD14*

gene is associated with asthma, perhaps providing some of the structural explanation for the hygiene hypothesis (63). This receptor may be part of the signaling mechanisms that mediate the proposed protective effects of childhood infections on asthma development.

Genes for inflammation and tissue damage

Tumor necrosis factor. Tumor necrosis factor (TNF) is an inflammatory cytokine found in increased concentrations in asthmatic airways (64) and in lavage fluid from asthmatic lungs (65). The TNF and lymphotoxin α and β genes are within the human major histocompatibility complex on chromosome 6p (66, 67). Constitutional variation in the level of secretion of TNF by peripheral blood lymphocytes or monocytes has been established in association with polymorphism in the TNF gene cluster and the *HLA-DRB1* locus (68–70).

Polymorphisms in the TNF genes have been associated with the presence of asthma (71). These polymorphisms act by enhancing the inflammatory process rather than modifying the IgE-mediated allergic response.

The high-affinity receptor for IgE (Fc ϵ RI) is the central trigger of the atopic response (64). It is multimeric, made up of one alpha, one beta, and two gamma chains. The receptor is also found in an alpha/gamma2 form. The alpha chain binds IgE, and the gamma chains carry out intracellular signaling. The beta chain is not necessary for receptor function but acts as an amplifying element.

Fc ϵ RI molecules are found on the surface of mast cells in the skin and mucosal lining of the airways and intestinal tract, and on basophils. The receptor binds circulating IgE molecules and holds them at the cell surface. The presence of antigens results in binding and cross-linking of IgE molecules. The resulting aggregation of receptors induces multiple signaling pathways that control diverse effector responses. These responses include secretion of allergic mediators and induction of cytokine gene expression, resulting in release of molecules such as *IL-4*, *IL-6*, TNF- α , and granulocyte-macrophage colony-stimulating factor. These responses are central to the induction and maintenance of allergic inflammation and may confer physiologic protection in parasite infections (64). Fc ϵ RI also is present on antigen-presenting cells, such as dendritic cells, and participates in IgE-mediated antigen presentation and processing (72).

Genes for asthma severity and bronchial hyperreactivity

Fc ϵ RI. The β chain of the high-affinity receptor for IgE (Fc ϵ RI) acts as an approximately sevenfold amplifying element of the receptor response to activation (65). Polymorphism within the chain has been associated with asthma (73), allergy (74), bronchial hyperresponsiveness (75), and atopic dermatitis (76). These variations seem to be associated with severe atopic disease.

***IL-4*.** Genetic markers within and around chromosome 5q31-33 have been linked to total serum IgE concentration in the United States (77) and Holland (78, 79). These markers

provide strong evidence of one or more loci in 5q31-33 closely involved in raised serum IgE levels and bronchial hyperresponsiveness. The gene encoding *IL-4* is within this region and is a possible candidate for the reported genetic linkage.

Because *IL-4* stimulation can influence mast cell responsiveness to IgE-mediated signaling, and because genetic variants in *IL-4* can modify *IL-4* gene transcription, these sequences also may modify asthma severity. In a study of 772 White and African-American asthmatics, the presence of the mutant *IL-4* promoter allele was associated with forced expiratory volume in 1 second of less than 50 percent among White study subjects. In Japan, polymorphisms of the *Ile50* allele of the *IL-4* receptor gene were associated with increased moderate-to-severe atopic asthma, especially in infants with onset within 2 years of age (80). Fatal and near-fatal asthma and severe airflow obstruction have been associated with polymorphisms in the *IL-4* 589T and *IL-4* RA576R alleles, respectively (81).

***IL-13*.** The Th2 cytokines *IL-4* and *IL-13* have a critical role in IgE synthesis. *IL-13* is closely related to *IL-4*, and the two genes arose from duplication of a single precursor. The receptors for both cytokines share an alpha chain (*IL-4R α*), which, when knocked out, greatly reduces IgE production (82). *IL-13* appears to affect asthmatic airways beyond simply enhancing IgE production (83).

β_2 Adrenergic receptor gene. Another candidate gene mapped to chromosome 5q is the β_2 adrenergic receptor (*AR*) gene. Several lines of evidence from experimental asthma have suggested that β_2 -AR may be related to asthma (84, 85). Two common polymorphisms of this gene at codons 16 and 27, *Arg16-Gly* and *Gln27-Glu*, were present with high allelic frequencies in both asthmatic and normal populations (86). Results from in vitro β_2 -AR experiments, using Chinese hamster fibroblast cells transfected with human airway smooth-muscle cells in tissue culture, suggested that these two polymorphisms are involved in agonist-promoted receptor down-regulation (87, 88). A substitution of arginine for glycine at codon 16 resulted in increased down-regulation of β_2 -AR after agonist challenge. In contrast, a substitution glutamic acid for glutamine at codon 27 conferred attenuated down-regulation. However, results from several small clinical studies were inconsistent, in part because these two polymorphisms were common and in linkage disequilibrium. The polymorphism at codon 16 was reported to be associated with asthma severity, nocturnal asthma, and airway hyperresponsiveness (86, 89, 90) but not with fatal or near-fatal asthma (91). On the contrary, Hall et al. (92) reported that *Gln27-Glu* was associated with lower airway reactivity in subjects with mild-to-moderate asthma.

***ADAM-33*.** The *ADAM-33* gene has recently been shown to be expressed by lung fibroblasts and bronchial smooth muscle cells (93), and it has been suggested that polymorphisms may influence smooth muscle cell and fibroblast proliferation, possibly leading to increased inflammation (94). Further work is needed to confirm the precise role of *ADAM-33* in asthma development.

PERINATAL RISK FACTORS

The greater influence of maternal compared with paternal asthma and atopy on the development of asthma in offspring suggests a role of the perinatal environment. In this section, we first review environmental risk factors occurring during the intrauterine period, which is often neglected even in studies that follow up children from birth. Factors that influence asthma development through allergic sensitization of the fetus or through alteration of the fetal environment are considered. Next, factors occurring in the neonatal period, including diet and lactation, and the nursery environment, are reviewed. Finally, we comment on environmental risk factors in early childhood.

Intrauterine risk factors for atopy and asthma development

Despite a large literature on risk factors for the development of asthma in children, almost nothing is known about the role of intrauterine factors (95). Aspects of the fetal environment that have been implicated in asthma development in the offspring include in utero immune responses and inadequate oxygenation and lung maturation.

Allergic sensitization in the fetus. Atopy is one of the most important risk factors for developing asthma, and nearly all asthmatics have altered immune responses (96). The human immune response begins in utero, and gestation and early childhood are thought to be the most influential periods with regard to atopic expression (97). Exposure to allergens begins during the perinatal period. The human fetus appears to produce IgE, but at relatively low levels. Nonetheless, higher total IgE levels in cord blood (>0.8 kU/liter) have been observed and were associated with increased risk for atopic disease in infancy, including atopic dermatitis (98, 99), and urticaria from food allergy (100), but the majority of studies have found little predictive value of cord blood IgE on asthma or asthmatic symptoms (98–107) because of the low sensitivity and predictive value of IgE (98, 102, 107–109). However, none of the studies of the predictive value of cord blood IgE has been conducted among subjects with any substantial understanding of their genetic polymorphisms, examples of which have just been described, or a detailed examination of their residential aeroallergen exposure, reviewed below.

It is widely assumed that maternal IgE does not cross the placenta (110), although IgG does (111), and that increased cord blood IgE may be due to intrauterine sensitization. Neonatal sensitivity to cow's milk, penicillin, helminthes, and grass pollen has been observed (112–118). Alternatively, elevated neonatal IgE may be due to nonspecific spontaneous IgE synthesis or to transplacental transfer of maternal IgE antibodies to cord blood that promotes fetal antibody formation (95). It has been shown that cord blood lymphocytes are stimulated by food and inhaled allergens (119, 120) but not known is whether this is a "sensitization" response, which predisposes the neonate to an allergic response, or normal immune system maturation (95). In one recent study of children with a family history of asthma,

elevated IgE levels at 6 months of age did predict asthma in the children at age 6 years (121).

Maternal transfer of IgG antibodies to the fetus confers immunity to some infectious diseases (111). Protection against other infectious diseases may itself lower atopy and asthma risk, since infections can alter mucosal barriers and influence immune responses (95). Maternal vaginitis and febrile infection during pregnancy was shown to increase asthma risk in offspring at age 7 years (122), as did infection of the amniotic cavity (123). It is also possible that maternal IgG response to some foods may protect infants from sensitization, but this possibility remains controversial (124–128). Vassella et al. (129) observed that elevated cord blood IgE and IgG were associated with fewer allergies during the first 18 months, particularly in infants with a family history of allergy.

Induction of neonatal IgE and IgG antibodies as a result of changing the maternal diet has been the subject of intervention trials (130, 131). Cow's milk, egg protein, fish, and peanuts were excluded from the maternal diet, whereas, in other studies, maternal milk and egg consumption were increased to stimulate IgG antibody development; however, neither strategy appeared successful (132). Because these trials examined only a main effect of diet, they do not inform us of any role that maternal Ig status owing to diet may have on infant allergy when considered in combination with other genetic and environmental risk factors.

A recent trial influenced the fetal environment of 132 high-risk infants; their mothers were randomly given the oral probiotic *Lactobacillus* GG during the later stages of pregnancy, and the newborns directly received the probiotic through breast milk after delivery (133). At age 2 years, children given the probiotic had half the rate of atopic eczema of the placebo group (23 percent vs. 46 percent, $p = 0.008$). Overall, six children developed asthma, who were not meaningfully distinguished by their probiotic therapy.

Influences of the fetal environment. An adequate fetal environment is critical to the growth and development of the fetus. Inadequate oxygenation and/or nutrition can lead to disruption of lung maturation. If early or severe enough, irreversible pulmonary abnormalities may persist (134). Maternal smoking during pregnancy is indisputably linked to fetal growth retardation, and passive smoke exposure may have similar effects (135–137). Maternal smoking also appears to increase the risk of asthma in the infant (138–144), although it has been difficult to disentangle the intrauterine risk from the effects of passive neonatal exposure (145). Recent work suggests that in utero exposure to maternal smoking without postpartum exposure to environmental tobacco smoke increases the risk of a child having physician-diagnosed asthma, although exposure to environmental tobacco smoke during childhood was related to wheeze but not asthma (146).

In addition to presumably conferring a genetic risk, maternal asthma, a condition associated with impaired respiratory function and possibly decreased oxygenation of the fetus, also may affect asthma development in offspring by affecting fetal development. This may explain why, in the genetic studies, maternal asthma confers more risk than paternal asthma (41, 42). Asthmatic status during pregnancy

has been linked in some studies but not others to poor birth outcomes. Results are equivocal regarding gestational diabetes (123, 147, 148), preterm labor (123, 147, 149–152), preterm delivery (147, 148, 150–154), fetal growth (155–157), and low birth weight (147, 156–161). A consistent, increased risk of hypertension during pregnancy among asthmatics has been found in prior research (123, 147, 148, 151, 153, 154, 156). Recently, umbilical artery flow velocity from Doppler ultrasound was found to be significantly reduced at 18 weeks' gestation in moderately and severely asthmatic mothers (162). Intrauterine exposure to beta agonists (163) and poorly managed maternal asthma (164, 165) have been associated with asthma development in children, but the independent effects of each have not been disentangled. These risk factors may contribute significantly to the burden of asthma, because 3.7–8.4 percent of US pregnancies have been estimated to be affected by maternal asthma (166).

Perinatal outcomes that indicate a compromise in the fetal environment have been inconclusively linked to asthma development. Preterm delivery has been shown to be possibly associated with asthma development in children (156, 167, 168), as has low birth weight (169–171). Very-low-birth-weight infants (<1,500 g) rarely have been studied, but, in one study, risks of asthma appeared to be associated with very low birth weight itself but not with other perinatal factors (172).

One Norwegian study tried to distinguish between maternal obstetric conditions (hyperemesis, hypertension, and preeclampsia) and uterine factors (antepartum hemorrhage, preterm contractions, placental insufficiency, and uterine growth restriction) and found an association only with the latter group for asthma at age 4 years. These results were observed in both atopic and nonatopic parents. IgE status of the children at birth was not recorded (173). In a recent Finnish study, risk of atopy until 31 years of age increased linearly with increasing gestational age from 35 weeks on, particularly among children of farmers, although farmers' children overall were at lower risk of atopy (174). We speculate that children born preterm and with immature immune systems react differently to allergens and antigens than do infants born later. In the Finnish study, no relations were observed between risk of atopy and asthma, indicating the complex etiology of that disease and the importance of also studying the genotype. The role of mother's education (11, 15, 18) and family income (15, 16, 22), which are associated with negative perinatal outcomes and increased asthma in children, also needs to be addressed in future studies.

Other perinatal factors implicated as possible risk factors for atopic disease and asthma development include increased ponderal index and decreasing maternal age (175), stress during pregnancy (176), delivery by cesarean section (177), hypertensive disorders of pregnancy and gestational diabetes (123), use of prostaglandins and hormones during pregnancy (178), breech and instrumental delivery (179), and early or threatened labor and malpresentation of the fetus (180). Male gender (2, 15, 16, 18, 181) is a well-recognized risk factor. Having older siblings was protective for asthma diagnosed after age 2 years in one recent study but was a risk

factor for earlier diagnosis (182), also pointing to the importance of deconstructing the risk factors for intrauterine, early neonatal, and later infancy exposure.

The influence of lactation and diet

The role of lactation in infant atopy and asthma remains controversial. Studies have reported protective effects of breastfeeding on asthma (183–185) and recurrent wheeze (16, 186, 187), increased risk of asthma related to breastfeeding (186, 188–190), and no relation between the two (16, 191, 192). Methodological issues in these studies include differences in defining breastfeeding, in study populations, and in age of the child when the outcome is assessed.

Breastfeeding has been variously defined as any versus none (16, 183, 187, 191), by duration (185, 186, 188), or as exclusive breastfeeding versus combined with formula or solid food (184, 189, 190, 192). None of these definitions consistently has been related to asthma outcome. For example, Oddy et al. reported that exclusive breastfeeding for 4 months or more was protective against asthma development (184). Takemura et al. reported a small increased risk of asthma among children exclusively breastfed (190). Wright et al. described a threefold increased risk of asthma among atopic children who were exclusively breastfed for 4 months or more and whose mothers had asthma (189). In another study, Wilson et al. reported that introduction of solid food before age 4 months increased the risk of wheeze, but exclusive breastfeeding was unrelated to asthma diagnosis (192).

Similarly, different study populations have provided quite different answers to the question of whether breastfeeding influences asthma development. Studies of population-based samples have yielded protective effects (183, 185), have suggested increased risk (188, 190), and have found no association (16, 192). Areas in which the prevalence of asthma is high have been found to have both a protective association of breastfeeding in Australia (184) and an increased risk in New Zealand (188), while a study of atopic families found no association (191).

There is somewhat more consistency when the literature is examined according to age of the child. Studies of young children (to age 6 years) often have reported that breastfeeding reduces the risk of asthma (183–185), while studies of older children (190) and young adults (188) are more likely to report that breastfeeding increases risk. In a large Italian study (186), breastfeeding for 6 months or more reduced the risk of wheeze in the first 2 years of life but increased the risk of late-onset wheeze. The Tucson Respiratory Study reported that breastfeeding prevented asthma when children were assessed at age 6 years (187), but breastfeeding increased the risk of asthma at age 13 years (189) among children whose mothers were asthmatic.

Any protective effect of lactation may be influenced by the mother's own immune status. Factors that suppress IgE have been reported in colostrum, the first-expressed breast milk (193). A randomized trial to compare cow's milk with banked human milk in preterm infants reported greater allergic disease in the infants exposed to cow's milk by 18 months of age (194). It is possible that exposure to some

food allergens in breast milk increases IgE sensitization (195) and that breast milk itself may increase milk-specific IgE (196), perhaps from the mother's own diet of cow's milk. Thus, maternal diet is an important factor when considering the effect of lactation on atopy (197).

Trials of both maternal and infant avoidance of allergenic foods have indicated a lower risk of food allergies but not asthma in the intervention group (130, 198). Only one trial has been known to combine food allergen avoidance with dust mite restriction, and, while reduced asthma and atopy were observed at 1 year of age (199), the effect was diminished at age 2 years (200). None of the trials of food avoidance considered gene-environment interactions; new trials should consider genetic susceptibility as an effect modifier.

Early neonatal risk factors

It recently was observed that, after adjustment for several potentially confounding factors, newborns who spent their first night in a communal nursery were at increased risk of developing hay fever (odds ratio = 1.48, 95 percent confidence interval: 1.23, 1.77) (201). The authors speculated that infants in the nursery were more likely to experience low-dose and short-duration exposure to nonfamilial microorganisms. This study was conducted in a 1970 birth cohort, and the care of newborns is very different today, particularly with the greater use of neonatal intensive care nurseries. However, these nurseries can themselves expose infants to a range of microorganisms that may be important for sensitization to asthma or to protection against future asthma (202). It also is known that the early extrauterine environment of the infant's immune system is dependent on gut microflora through lipopolysaccharides in Gram-negative bacteria and that different bacteria induce various cytokines (203, 204). This knowledge raises the possibility that use of antibiotics in the neonatal nursery may enhance the future development of asthma. Older children treated with antibiotics have shown increased rates of asthma (205). Antibiotic use in early infancy has been observed to strongly increase risk of asthma at ages 5–10 years (odds ratio = 4.0, 95 percent confidence interval: 1.6, 10.6) (206). These findings also raise hypotheses about the use of antibiotics in the newborn intensive care nursery as well as obstetric antibiotic use, although confounding by disease indication in future studies will require careful control.

It has been observed that babies born preterm, of low birth weight, and needing positive pressure ventilation at birth, as well as having mothers who smoked during pregnancy, have an increased susceptibility to asthma (207–209). Babies with low 1- and 5-minute Apgar scores are also at increased risk (179), but all of these factors are more common in those who have spent time in the neonatal intensive care nursery. Consequently, the independent effects of antenatal versus early postnatal exposures have not been satisfactorily delineated, perhaps because, to our knowledge, they have never been studied in the same cohort.

Environmental risk factors in infancy and early childhood

Allergens in house dust have been controversially implicated in the development and severity of asthma in children. House dust mite (210–216), cat (217–222), dog (223–225), cockroach (218, 226–229), and fungi (211, 213–216, 230–234) allergens are among the most important because of their suspected role in the development and exacerbation of asthma. Recent cohort studies that measured allergen exposure in infancy and followed children for recurrent wheeze and asthma have failed to find a strong association with dust mite or cat allergen (191, 235–237). One study that stratified by maternal asthma status suggested an interaction between exposure and genetic susceptibility (237).

At least three studies (35, 238, 239) found an association between mold in homes and respiratory symptoms. Recent studies have reported that house dust endotoxins independently increased symptoms of wheeze in the first year of life (240), and in utero allergen sensitization is common and may involve *IL-5* regulation (241). Endotoxin, a lipopolysaccharide from the outside layer of Gram-negative bacteria found in the fecal material of large mammals, is of particular interest for the hygiene hypothesis. This hypothesis followed observations that children reared on farms had lower rates of asthma than did children from urban environments (242). A recent publication (243) found that endotoxin levels from the mattresses of children aged 6–13 years in rural European communities, but not necessarily from farms, were inversely related to atopic asthma and hay fever as well as to cytokine production by leukocytes, indicating increased tolerance of exposure to endotoxin and other allergens.

A particularly important indoor air contaminant in the exacerbation of asthma, environmental tobacco smoke, is associated with a wide range of acute and chronic effects (142, 244–246.) The US Environmental Protection Agency calculated that the children of mothers who smoked 10 or more cigarettes daily had a 90–600 percent increased risk of developing asthma and that 48–83 percent of all cases of asthma in the children of these women could be attributed to their mother's smoking (247). A recent California study reported that maternal smoking primarily increased the risk of early-onset persistent asthma, particularly in children with a history of parental asthma, suggesting the importance of further defining the genotype of at-risk children (248).

Limited human and animal data suggest that air contaminants, particularly ozone and nitrogen dioxide, may modulate immune responses to inhaled biologicals (139, 199, 249, 250). Viral (251–253) and respiratory (44, 254, 255) infections before the age of 2 years have been associated with increased risk of asthma. Indoor heating sources (34, 256), use of gas stoves (34, 257, 258), and nitrogen dioxide exposure (258–263) have all been related to increased infant respiratory symptoms.

SUMMARY AND THEORETICAL MODEL

Asthma development and severity are almost certainly influenced by a multifactorial etiology that includes genetic, immunologic, and environmental factors. Many of these risk




FIGURE 1. Hypothesized associations of genetic, intrauterine, and perinatal risk factors with asthma onset and severity in infancy, modified from Wahn (267). Ig, immunoglobulin.

factors may start to exert their effect during intrauterine life (264–266). Future research, by building on these risk factors and a detailed investigation of the mother's own asthma status during the index pregnancy, may start to shed some light on these relations.

No single factor is likely to be responsible for the observed increase in asthma onset and severity, particularly the increase in minority populations. It is almost certainly due to new environmental factors, since the gene pool is unlikely to have changed meaningfully in the last 20–30 years. New environmental factors will influence phenotypes genetically susceptible to them, at selected stages in the development of disease. We have suggested that asthma development may start as early as the fetal period and is certainly well established during infancy.

The hypothetical roles of the major groups of risk factors, discussed throughout this review, are shown in figure 1, which has been modified from Wahn (267) to emphasize the role of intrauterine and early neonatal exposures. It is likely that a complex interplay of these factors is responsible for asthma development and severity. Current knowledge suggests that development of asthma requires polymorphisms in several genes in the same person, which will lead to complex gene-gene and gene-environment interactions. These interactions are likely to come into play at different stages during early life, as the child becomes exposed to a variety of environmental insults. These complex interactions make it difficult to initially identify candidate genes and, once identified, to understand the full role they play in asthma development and severity.

Although a number of susceptibility genes for asthma have been identified, it is almost certain that the majority remain to be described. Analysis of shared haplotypes (linear arrangements of alleles on a chromosome) of affected persons from "founder populations" with a common ancestor is increasingly being used to identify candidate asthma genes (268). Furthermore, with publication of the human genome, candidate asthma genes are likely to be identified at an increasing rate. The expected plethora of candidate genes will pose problems of its own unless fundamental epidemiologic principles are followed. These principles include the need for replicating findings in alternative

populations, considering publication bias in the evolving literature, systematically reviewing evidence for a candidate gene, and conducting a meta-analysis of its effects when appropriate. Application of other criteria for causality, well known to epidemiologists, will be important. Included are strong associations unconfounded by population stratification, biologic plausibility (what is the function of the gene's protein products?), and dose response based on homo- or heterozygosity. To organize this complex body of data into testable hypotheses, increasingly sophisticated theoretical modeling of new associations will be required.

Although exposure to perinatal factors, allergens, and other environmental risk factors increases the risk of asthma, only a portion of exposed neonates will develop asthma. We conclude that neonates' susceptibility will vary based on their genetic and immunologic predisposition, and identification of at-risk genotypes will permit more precise specification of environmental risk models. Study cohorts are required in which both parents and the index child can be genotyped and the child prospectively followed from *early pregnancy* through infancy. These studies will enable analysis of the gene-gene and gene-immunologic-environment interactions likely to fundamentally influence asthma onset and severity. Our model has utility in considering the interplay of the genotype with environmental factors, because it points to the focus of clinical specialists: obstetricians, perinatologists, and pediatricians. In the future, the management of patients is likely to be influenced by pharmacogenetic studies of the responsiveness of patients with selected polymorphisms to a particular medication. The model also emphasizes the need for continuity of care among patients at risk of asthma and the importance of understanding the complex nature of asthma etiology if more effective prevention programs are to be launched. Knowing the genetic risk of patients may permit more targeted allergen avoidance strategies (269).

The model is also overly simple. A particular polymorphism may influence the course of asthma at more than one stage. As described earlier, *IL-13* polymorphisms have already been shown to increase the risk of atopy and asthma severity. At the same time, environmental interactions are becoming more apparent. Babies born preterm and also

exposed to environmental tobacco smoke are at an enhanced risk of asthma. There are extremely difficult technical problems, which are outside the scope of this review, in designing studies able to identify gene-gene-environment-environment interactions, and we are aware of none conducted to date for any disease. However, asthma is a disease for which such a complex array of factors will almost certainly be needed to explain substantial amounts of variance in asthma risk.

ACKNOWLEDGMENTS

This work was supported by grants AI41040, ES07456, and ES05410 from the National Institutes of Health, Bethesda, Maryland.

The authors are particularly grateful to Lia Kidd for her expertise in preparing the manuscript for publication.

REFERENCES

1. Woolcock AJ, Peat JK. Evidence for the increase in asthma worldwide. *Ciba Found Symp* 1997;206:122–34.
2. Mannino DM, Homa DM, Pertowski CA, et al. Surveillance for asthma—United States, 1960–1995. *Mor Mortal Wkly Rep CDC Surveill Summ* 1998;47:1–28.
3. Rappaport S, Boodram B. Forecasted state-specific estimates of self-reported asthma prevalence—United States, 1998. *MMWR Morb Mortal Wkly Rep* 1998;47:1022–5.
4. Cherry DK, Woodwell DA. National Ambulatory Medical Care Survey: 2000 summary. Advance data from Vital and Health Statistics, no. 328. Hyattsville, MD: National Center for Health Statistics, 2001.
5. McCaig LF, Ly N. National Hospital Ambulatory Medical Care Survey: 2000 Emergency Department summary. Advance data from Vital and Health Statistics, no. 326. Hyattsville, MD: National Center for Health Statistics, 2001.
6. Beckett WS, Belanger K, Gent JF, et al. Asthma among Puerto Rican Hispanics: a multi-ethnic comparison study of risk factors. *Am J Respir Crit Care Med* 1996;154:894–9.
7. Persky VW, Slezak J, Contreras A, et al. Relationship of race and socioeconomic status with prevalence, severity and symptoms of asthma in Chicago school children. *Ann Allergy Asthma Immunol* 1998;81:226–71.
8. Gergen PJ. Race, income, urbanicity, and asthma hospitalization in California: a small area analysis. *Chest* 1998;113:1277–84.
9. Goodman DC, Stukel TA, Chang CH. Trends in pediatric asthma hospitalization rates: regional and socioeconomic differences. *Pediatrics* 1998;101:208–13.
10. Lara M, Morgenstern H, Daun N, et al. Elevated asthma morbidity in Puerto Rican children: a review of possible risk and prognostic factors. *West J Med* 1999;170:75–84.
11. Leidy NK, Coughlin C. Psychometric performance of the Asthma Quality of Life Questionnaire in a US sample. *Qual Life Res* 1998;7:127–34.
12. Coultas DB, Gong H, Grad R, et al. Respiratory disease in minorities of the United States. *Am J Respir Crit Care Med* 1993;149:S93–S131.
13. Croner S, Kjellman N. Natural history of bronchial asthma in childhood. *Allergy* 1992;47:150–7.
14. Seigel SC, Rachelefsky GS. Asthma in infants and children. Part I. *J Allergy Clin Immunol* 1985;76:1–15.
15. Weitzman M, Gortmaker SL, Sobol AM, et al. Recent trends in the prevalence and severity of childhood asthma. *JAMA* 1992;268:2673–7.
16. Schwartz J, Gold D, Dockery DW, et al. Predictors of asthma and persistent wheeze in a national sample of children in the United States. Association with social class, perinatal events, and race. *Am Rev Respir Dis* 1990;142:555–62.
17. Gold DR, Rotnitzky A, Damokosh AI, et al. Race and gender differences in respiratory illness prevalence and their relationship to environmental exposures in children 7–14 years of age. *Am Rev Respir Dis* 1993;148:10–18.
18. Weitzman M, Gortmaker S, Sobol A. Racial, social, and environmental risks for childhood asthma. *Am J Dis Child* 1990;144:1189–94.
19. Graves EJ. National hospital discharge survey: annual summary, 1987. *Vital Health Stat* 13 1989;Apr(99):1–60. (Publication (PHS) 89-1760. PB90-158858. PC A04 MF A01).
20. Halfon N, Newacheck PW. Trends in the hospitalization for acute childhood asthma, 1970–84. *Am J Public Health* 1986;76:1308–11.
21. Evans R III. Asthma among minority children. *Chest* 1992;101:368S–71S.
22. Carr W, Zeitel L, Weiss K. Variations in asthma hospitalizations and deaths in New York City. *Am J Public Health* 1992;82:59–65.
23. Wissow LS, Gittelsohn AM, Szklo M, et al. Poverty, race and hospitalization for childhood asthma. *Am J Public Health* 1988;78:777–82.
24. Gergen PJ, Mullally DI, Evans R III. Changing patterns of asthma hospitalization among children: 1979–1987. *JAMA* 1990;264:1688–92.
25. Weiss KB, Wagener DK. Asthma surveillance in the United States: a review of current trends and knowledge gaps. *Chest* 1990;95:179S–84S.
26. Weiss KB, Wagener DK. Changing patterns of asthma mortality: identifying target populations at high risk. *JAMA* 1990;264:1683–7.
27. Carter-Pokras OD, Gergen PJ. Reported asthma among Puerto Rican, Mexican-American, and Cuban children, 1982 through 1984. *Am J Public Health* 1993;83:580–2.
28. Burney PG, Chinn S, Rona RJ. Has the prevalence of asthma increased in children? Evidence from the national study of health and growth 1973–1986. *BMJ* 1990;300:1306–10.
29. Rona RJ, Chinn S, Burney PG. Trends in the prevalence of asthma in Scottish and English primary school children 1982–1992. *Thorax* 1995;50:992–3.
30. Anthracopoulos M, Karatza A, Liolios E, et al. Prevalence of asthma among schoolchildren in Patras, Greece: three surveys over 20 years. *Thorax* 2001;56:569–71.
31. Venn A, Lewis S, Cooper M, et al. Increasing prevalence of wheeze and asthma in Nottingham primary schoolchildren 1988–1995. *Eur Respir J* 1998;11:1324–8.
32. Ferguson AL. Diagnosis in children. In: Fitzgerald JM, Ernst P, Boulet LP, et al, eds. Evidence based asthma management. Hamilton, Ontario, Canada: BC Decker Inc, 2001.
33. Martinez FD, Wright AL, Taussig LM, et al. Asthma and wheezing in the first six years of life. *N Engl J Med* 1995;332:133–8.
34. Triche E, Belanger K, Beckett W, et al. Infant respiratory symptoms associated with indoor heating sources. *Am J Respir Crit Care Med* 2002;166:1105–11.
35. Gent JF, Ren P, Belanger K, et al. Levels of household mold associated with respiratory symptoms in the first year of life

- in a cohort at risk for asthma. *Environ Health Perspect* 2002; 110:A781–6.
36. Ng Man Kwong G, Proctor A, Billings C, et al. Increasing prevalence of asthma diagnosis and symptoms in children is confined to mild symptoms. *Thorax* 2001;56:312–14.
 37. Bleecker ER, Postma DS, Meyers DA. Evidence for multiple genetic susceptibility loci for asthma. *Am J Respir Crit Care Med* 1997;156:S113–16.
 38. Marsh DG, Meyers DA. A major gene for allergy—fact or fancy? *Nat Genet* 1992;2:252–4.
 39. Sandford A, Weir T, Pare P. The genetics of asthma. *Am J Respir Crit Care Med* 1996;153:1749–65.
 40. Aberg A. Familial occurrence of atopic disease: genetic versus environmental factors. *Clin Exp Allergy* 1993;23:829–34.
 41. Litonjua AA, Carey VJ, Burge HA, et al. Parental history and the risk for childhood asthma. Does mother confer more risk than father? *Am J Respir Crit Care Med* 1998;158:176–81.
 42. Holberg CJ, Morgan WJ, Wright AL, et al. Differences in familial segregation of FEV₁ between asthmatic and nonasthmatic families. Role of a maternal component. *Am J Respir Crit Care Med* 1998;158:162–9.
 43. Edfors-Lubs ML. Allergy in 7,000 twin pairs. *Acta Allergol* 1971;26:249–85.
 44. Sherman CB, Tosteson TD, Tager IB, et al. Early childhood predictors of asthma. *Am J Epidemiol* 1990;132:83–95.
 45. Räsänen M, Kaprio J, Laitinen T, et al. Perinatal risk factors for asthma in Finnish adolescent twins. *Thorax* 2000;55:25–31.
 46. Laitinen T, Räsänen M, Kaprio J, et al. Importance of genetic factors in adolescent asthma: a population-based twin-family study. *Am J Respir Crit Care Med* 1998;157:1073–8.
 47. Räsänen M, Laitinen T, Kaprio J, et al. Hay fever—a Finnish nationwide study of adolescent twins and their parents. *Allergy* 1998;53:885–90.
 48. Räsänen M, Kaprio J, Laitinen T, et al. Perinatal risk factors for hay fever—a study among 2550 Finnish twin families. *Twin Res* 2001;4:392–9.
 49. Los H, Postmus PE, Boomsma DI. Asthma genetics and intermediate phenotypes: a review from twin studies. *Twin Res* 2001;4:81–93.
 50. Strachan DP, Wong HJ, Spector TD. Concordance and interrelationship of atopic diseases and markers of allergic sensitization among adult female twins. *J Allergy Clin Immunol* 2001;108:901–7.
 51. Lander ES, Schork NJ. Genetic dissection of complex traits. *Science* 1994;265:2037–48.
 52. A genome-wide search for asthma susceptibility loci in ethnically diverse populations. The Collaborative Study on the Genetics of Asthma (CSGA). *Nat Genet* 1997;15:389–92.
 53. Daniels SE, Bhattacharaya S, James A, et al. A genome-wide search for quantitative trait loci underlying asthma. *Nature* 1996;383:247–50.
 54. Cookson WO. Asthma genetics. *Chest* 2002;121:7S–13S.
 55. Borish L, Mascali JJ, Klinnert M, et al. SSC polymorphisms in interleukin genes. (Erratum). *Hum Mol Genet* 1995;4:974.
 56. Noguchi E, Shibasaki M, Arinami T, et al. Association of asthma and the interleukin-4 promoter gene in Japanese. *Clin Exp Allergy* 1998;28:449–53.
 57. Walley AJ, Cookson WOCM. Investigation of an interleukin-4 promoter polymorphism for associations with asthma and atopy. *J Med Genet* 1996;33:689–92.
 58. Graves PE, Kabesch M, Halonen M, et al. A cluster of seven tightly linked polymorphisms in the *IL-13* gene is associated with total serum IgE levels in three populations of white children. *J Allergy Clin Immunol* 2000;105:506–13.
 59. Kruse S, Japha T, Tedner M, et al. The polymorphisms *S503P* and *Q576R* in the interleukin-4 receptor alpha gene are associated with atopy and influence the signal transduction. *Immunology* 1999;96:365–71.
 60. Kruse S, Forster J, Kuehr J, et al. Characterization of the membrane-bound and a soluble form of human *IL-4* receptor alpha produced by alternative splicing. *Int Immunol* 1999;11:1965–70.
 61. Hackstein H, Hecker M, Kruse S, et al. A novel polymorphism in the 5' promoter region of the human interleukin-4 receptor alpha-chain gene is associated with decreased soluble interleukin-4 receptor protein levels. *Immunogenetics* 2001;53:264–9.
 62. Howard TD, Koppelman GH, Xu J, et al. Gene-gene interaction in asthma: *IL4RA* and *IL13* in a Dutch population with asthma. *Am J Hum Genet* 2002;70:230–6.
 63. Baldini M, Lohman IC, Halonen M, et al. A Polymorphism* in the 5' flanking region of the *CD14* gene is associated with circulating soluble *CD14* levels and with total serum immunoglobulin E. *Am J Respir Cell Mol Biol* 1999;20:976–83.
 64. Broide DH, Lotz M, Cuomo AJ, et al. Cytokines in symptomatic asthma airways. *J Allergy Clin Immunol* 1992;89:958–67.
 65. Virchow JC Jr, Walker C, Hafner D, et al. T cells and cytokines in bronchoalveolar lavage fluid after segmental allergen provocation in atopic asthma. *Am J Respir Crit Care Med* 1995;151:960–8.
 66. Dunham I, Sargent CA, Trowsdale J, et al. Molecular mapping of the human major histocompatibility complex by pulsed-field gel electrophoresis. *Proc Natl Acad Sci U S A* 1987;84:7237–41.
 67. Nedospasov SA, Shakhov AN, Turetskaia RL, et al. Molecular cloning of human genes coding tumor necrosis factors: tandem arrangement of alpha- and beta-genes in a short segment (6 thousand nucleotide pairs) of human genome. (In Russian). *Dokl Akad Nauk SSSR* 1985;285:1487–90.
 68. Wilson AG, Symons JA, McDowell TL, et al. Effects of a tumour necrosis factor (TNF alpha) promoter base transition on transcriptional activity. *Proc Natl Acad Sci U S A* 1997;94:3195–9.
 69. Messer G, Spengler U, Jung MC, et al. Polymorphic structure of the tumor necrosis factor (TNF) locus: an *NcoI* polymorphism in the first intron of the human TNF-beta gene correlates with a variant amino acid in position 26 and a reduced level of TNF-beta production. *J Exp Med* 1991;173:209–19.
 70. Jacob CO, Fronek Z, Lewis GD, et al. Heritable major histocompatibility complex class II-associated differences in production of tumor necrosis factor alpha: relevance to genetic predisposition to systemic lupus erythematosus. *Proc Natl Acad Sci U S A* 1990;87:1233–7.
 71. Moffatt MF, Cookson WO. Tumour necrosis factor haplotypes and asthma. *Hum Mol Genet* 1997;6:551–4.
 72. Turner H, Kinet JP. Signalling through high-affinity IgE receptor Fc epsilonRI. *Nature* 1999;401:B24–30.
 73. Shirakawa T, Mao XQ, Sasaki S, et al. Association between atopic asthma and a coding variant of Fc epsilon RI beta in a Japanese population. *Hum Mol Genet* 1996;5:1129–30.
 74. Hill MR, James AL, Faux JA, et al. FcεRI-β polymorphism and risk of atopy in a general population sample. *BMJ* 1995; 311:776–9.
 75. van Herwerden L, Harrap SB, Wong ZY, et al. Linkage of high-affinity IgE receptor gene with bronchial hyperreactivity, even in absence of atopy. *Lancet* 1995;346:1262–5.
 76. Cox HE, Moffatt MF, Faux JA, et al. Association of atopic dermatitis to the beta subunit of the high affinity immu-

- noglobulin E receptor. (See comments). *Br J Dermatol* 1998; 138:182–7.
77. Marsh DG, Neely JD, Breazeale DR, et al. Linkage analysis of *IL4* and other chromosome 5q31.1 markers and total serum IgE concentrations. *Science* 1994;264:1152–6.
 78. Postma DS, Bleecker ER, Amelung PJ, et al. Genetic susceptibility to asthma—bronchial hyperresponsiveness coinherited with a major gene for atopy. *N Engl J Med* 1995;333: 894–900.
 79. Xu J, Levitt RC, Panhuysen CI, et al. Evidence for two unlinked loci regulating total serum IgE levels. *Am J Hum Genet* 1995;57:425–30.
 80. Takabayashi A, Ihara K, Sasaki Y, et al. Childhood atopic asthma: positive association with a polymorphism of *IL-4* receptor alpha gene but not with that of *IL-4* promoter or Fc epsilon receptor I beta gene. *Exp Clin Immunogenet* 2000;17: 63–70.
 81. Sandford AJ, Chagani T, Zhu S, et al. Polymorphisms in the *IL4*, *IL4RA* and *FCER1B* genes and asthma severity. *J Allergy Clin Immunol* 2000;106:135–40.
 82. Grunewald SM, Werthmann A, Schnarr B, et al. An antagonistic *IL-4* mutant prevents type I allergy in the mouse: inhibition of the *IL-4/IL-13* receptor system completely abrogates humoral immune response to allergen and development of allergic symptoms in vivo. *J Immunol* 1998;160:4004–9.
 83. Wills-Karp M, Luyimbazi J, Xu X, et al. Interleukin-13: central mediator of allergic asthma. *Science* 1998;282:2258–61.
 84. Barnes PJ, Dollery CT, MacDermot J. Increased pulmonary alpha-adrenergic and reduced beta-adrenergic receptors in experimental asthma. *Nature* 1980;285:569–71.
 85. Gatto C, Green TP, Johnson MG, et al. Localization of quantitative changes in pulmonary beta-receptors in ovalbumin-sensitized guinea pigs. *Am Rev Respir Dis* 1987;136:150–4.
 86. Reihnsaus E, Innis M, MacIntyre N, et al. Mutations in the gene encoding for the beta 2-adrenergic receptor in normal and asthmatic subjects. *Am J Respir Cell Mol Biol* 1993;8:334–9.
 87. Green SA, Turki J, Innis M, et al. Amino-terminal polymorphisms of the human beta 2-adrenergic receptor impart distinct agonist-promoted regulatory properties. *Biochemistry* 1994;33:9414–19. (Erratum published in *Biochemistry* 1994; 33:14368).
 88. Green SA, Turki J, Bejarano P, et al. Influence of beta 2-adrenergic receptor genotypes on signal transduction in human airway smooth-muscle cells. *Am J Respir Cell Mol Biol* 1995;13:25–33.
 89. Turki J, Pak J, Green SA, et al. Genetic polymorphisms of the beta 2-adrenergic receptor in nocturnal and non-nocturnal asthma. Evidence that Gly-916 correlates with the nocturnal phenotype. *J Clin Invest* 1995;95:1635–41.
 90. Martinez FD, Graves PE, Baldini M, et al. Association between genetic polymorphisms of the beta2-adrenoceptor and response to albuterol in children with and without a history of wheezing. *J Clin Invest* 1997;100:3184–8.
 91. Weir TD, Malleck N, Sandford AJ, et al. Genetic polymorphisms of the beta 2-adrenergic receptor in fatal and near-fatal asthma. (Abstract). *Am J Respir Crit Care Med* 1997;155: A257.
 92. Hall IP, Wheatley A, Wilding P, et al. Association of Glu 27 beta 2-adrenoceptor polymorphism with lower airway reactivity in asthmatic subjects. *Lancet* 1995;345:1213–14.
 93. Van Eerdewegh P, Little RD, Dupuis J, et al. Association of the *ADAM33* gene with asthma and bronchial hyperresponsiveness. *Nature* 2002;418:426–30.
 94. Shapiro SD, Owen CA. Clinical implications of basic research: *ADAM-33* surfaces as an asthma gene. *N Engl J Med* 2002;347:936–8.
 95. Bjorksten B, Kjellman NIM. Risk factors in the development of allergy. In: Schatz M, Zeiger R, Claman H, eds. *Asthma and immunological diseases in pregnancy and early infancy*. New York, NY: Marcel Dekker, 1998.
 96. Donovan CE, Finn PW. Immune mechanisms of childhood asthma. *Thorax* 1999;54:938–46.
 97. Peden DB. Development of atopy and asthma: candidate environmental influences and important periods of exposure. *Environ Health Perspect* 2000;108:475–82.
 98. Edenharter G, Bergmann RL, Bergmann KE, et al. Cord blood—IgE as risk factor and predictor for atopic diseases. *Clin Exp Allergy* 1998;28:671–8.
 99. Bergmann RL, Edenharter G, Bergmann KE, et al. Predictability of early atopy by cord blood IgE and parental history. *Clin Exp Allergy* 1997;27:752–60.
 100. Kaan A, Dimich-Ward H, Manfreda J, et al. Cord blood IgE: its determinants and prediction of development of asthma and other allergic disorders at 12 months. *Ann Allergy Asthma Immunol* 2000;84:37–42.
 101. Kjellmann NIM. IgE determinations in neonates is not suitable for general screening. *Pediatr Allergy Immunol* 1994;5:1–4.
 102. Eiriksson TH, Sigurgeirsson B, Ardal B, et al. Cord blood IgE levels are influenced by gestational age but do not predict allergic manifestations in infants. *Pediatr Allergy Immunol* 1994;5:5–10.
 103. Tariq SM, Arshad SH, Matthews SM, et al. Elevated cord serum IgE increases the risk of aeroallergen sensitization without increasing respiratory allergic symptoms in early childhood. *Clin Exp Allergy* 1999;29:1042–8.
 104. Halonen M, Stern D, Taussig LM, et al. The predictive relationship between serum IgE levels at birth and subsequent incidences of lower respiratory illnesses and eczema in infants. *Am Rev Respir Dis* 1992;146:866–70.
 105. Hide DW, Arshad SH, Twiselton R, et al. Cord serum IgE: an insensitive method for prediction of atopy. *Clin Exp Allergy* 1991;21:739–43.
 106. Hansen LG, Halken S, Host A, et al. Prediction of allergy from family history and cord blood IgE levels. A follow-up at the age of 5 years. Cord blood IgE IV. *Pediatr Allergy Immunol* 1993;4:34–40.
 107. Hansen LG, Host A, Halken S, et al. Cord blood IgE II. Prediction of atopic disease. A follow-up at the age of 18 months. *Allergy* 1992;47:397–403.
 108. Oldak E. Cord blood IgE levels as a predictive value of the atopic disease in early infancy: a review article. *Rocz Akad Med Białymst* 1997;42:13–17.
 109. Ruiz RG, Richards D, Kemeny DM, et al. Neonatal IgE: a poor screen for atopic disease. *Clin Exp Allergy* 1991;21: 467–72.
 110. Madani G, Heiner DC. Antibody transmission from mother to fetus. *Curr Opin Immunol* 1989;1:1157–64.
 111. Bramwell F. The transmission of antibodies. In: Bramwell F, ed. *The transmission of immunity from mother to young*. Amsterdam, the Netherlands: North-Holland, 1970:242–50.
 112. Businco L, Marchetti F, Pellegrini G, et al. Predictive value of cord blood IgE levels in “at risk” newborn babies and influence of type of feeding. *Clin Allergy* 1983;13:503–8.
 113. Delespesse G, Sarfati M, Lang G, et al. Prenatal and neonatal synthesis of IgE. *Monogr Allergy* 1983;18:83–95.
 114. Kimpen J, Callaert H, Embrechts P, et al. Influence of sex and gestational age on cord blood IgE. *Acta Paediatr Scand* 1989; 78:233–8.
 115. Host A, Halken S. A prospective study of cow milk allergy in Danish infants during the first 3 years of life. *Allergy* 1990; 45:587–96.

116. Host A, Husby S, Gjesing B, et al. Prospective estimation of IgG, IgG subclass and IgE antibodies to dietary proteins in infants with cow milk allergy. Levels of antibodies to whole milk protein, BLG and ovalbumin in relation to repeated milk challenge and clinical course of cow milk allergy. *Allergy* 1992;47:218–29.
117. Levin S, Altman Y, Sela M. Penicillin and dinitrophenyl antibodies in newborn and mothers detected with chemically modified bacteriophage. *Pediatr Res* 1971;5:87–8.
118. Weil GJ, Hussain R, Kumaraswami V, et al. Prenatal allergic sensitization to helminth antigen in offspring of parasite-infected mothers. *J Clin Invest* 1983;71:1124–9.
119. Warner JA, Miles EA, Jones AC, et al. Is deficiency of interferon gamma production by allergen triggered cord blood cells a predictor of atopic eczema? *Clin Exp Allergy* 1994;24:423–30.
120. Piccinni MP, Mecacci F, Sampognaro S, et al. Aeroallergen sensitization can occur during fetal life. *Int Arch Allergy Immunol* 1993;102:301–30.
121. Klinnert MD, Nelson HS, Price MR, et al. Onset and persistence of childhood asthma: predictors from infancy. *Pediatrics* 2001;108:E69.
122. Xu B, Pekkanen J, Jarvelin MR, et al. Maternal infections in pregnancy and the development of asthma among offspring. *Int J Epidemiol* 1999;28:723–7.
123. Wen SW, Demissie K, Liu S. Adverse outcomes in pregnancies of asthmatic women: results from a Canadian population. *Ann Epidemiol* 2001;11:7–12.
124. Lilja G, Dannaes A, Falth-Magnusson K, et al. Immune response of the atopic woman and foetus: effects of high- and low-dose food allergen intake during late pregnancy. *Clin Allergy* 1988;18:113–42.
125. Falth-Magnusson K, Oman H, Kjellman NI. Maternal abstinence from cow milk and egg in allergy risk pregnancies. Effect on antibody production in the mother and the newborn. *Allergy* 1987;42:64–73.
126. Dannaes A, Johansson SG. Clinical and immunological aspects of food allergy in childhood. II. Development of allergic symptoms and humoral immune response to foods in infants of atopic mothers during the first 24 months of life. *Acta Paediatr Scand* 1978;67:497–504.
127. Casimir G, Duchateau J, Gossart B, et al. Antibody against betalactoglobulin (IgG) and cow's milk allergy. (Abstract). *J Allergy Clin Immunol* 1985;75:206.
128. Lilja G, Dannaes A, Foucard T, et al. Effects of maternal diet during late pregnancy and lactation on the development of atopic diseases in infants up to eighteen months of age: in-vivo results. *Clin Exp Allergy* 1989;19:473–9.
129. Vassella CC, Odelram H, Kjellman NI, et al. High anti-IgE levels at birth are associated with a reduced allergy prevalence in infants at risk: a prospective study. *Clin Exp Allergy* 1994;24:771–7.
130. Falth-Magnusson K, Kjellmann NIM. Allergy prevention by maternal elimination diet during late pregnancy—a 5 year follow-up of randomized study. *J Allergy Clin Immunol* 1992;89:709–13.
131. Zeiger RS, Heller S, Mellon MH, et al. Genetic and environmental factors affecting the development of atopy through age 4 in children of atopic parents: a prospective randomized study of food allergen avoidance. *Pediatr Allergy Immunol* 1992;3:110–27.
132. Zeiger RS. Prevention of allergic disease in infancy. In: Schatz M, Zeiger R, Claman H, eds. *Asthma and immunological diseases in pregnancy and early infancy*. New York, NY: Marcel Dekker, Inc, 1998:761–811.
133. Kalliomäki M, Salminen S, Arvilommi H, et al. Probiotics in primary prevention of atopic disease: a randomised placebo-controlled trial. *Lancet* 2001;357:1076–9.
134. Horowitz S, Davis JM. Lung injury when development is interrupted by premature birth. In: McDonald J, ed. *Lung growth and development*. New York, NY: Marcel Dekker, Inc, 1997:577–610.
135. Martin TR, Bracken MB. Association of low birth weight with passive smoke exposure in pregnancy. *Am J Epidemiol* 1986;124:633–42.
136. Sadler L, Belanger K, Saftlas A, et al. Environmental tobacco smoke exposure and small-for-gestational-age birth. *Am J Epidemiol* 1999;150:695–705.
137. US Department of Health and Human Services. Women and smoking: a report of the Surgeon General—2001. Washington, DC: Centers for Disease Control and Prevention, National Center for Chronic Disease Prevention and Health Promotion, Office on Smoking and Health, 2002:272–368. (ISBN 0-16-050751-0).
138. Weitzman M, Gortmaker S, Walker DK, et al. Maternal smoking and childhood asthma. *Pediatrics* 1990;85:505–11.
139. Ehrlich R, Kattan M, Godbold J, et al. Childhood asthma and passive smoking. Urinary cotinine as a biomarker of exposure. *Am Rev Respir Dis* 1992;145:594–9.
140. Taylor B, Wadsworth J. Maternal smoking during pregnancy and lower respiratory tract illness in early life. *Arch Dis Child* 1987;62:786–91.
141. Li YF, Gilliland FD, Berhane K, et al. Effects of in utero and environmental tobacco smoke exposure on lung function in boys and girls with and without asthma. *Am J Respir Crit Care Med* 2000;162:2097–104.
142. Joad JP. Smoking and pediatric respiratory health. *Clin Chest Med* 2000;21:37–46.
143. Strachan DP, Cook DG. Health effects of passive smoking—5: parental smoking and allergic sensitisation in children. *Thorax* 1998;53:117–23.
144. Lodrup Carlsen KC, Carlsen KH. Effects of maternal and early tobacco exposure on the development of asthma and airway hyperreactivity. *Curr Opin Allergy Clin Immunol* 2001;1:139–43.
145. Lux AL, Henderson AJ, Pocock SJ. Wheeze associated with prenatal tobacco smoke exposure: a prospective longitudinal study. ALSPAC study team. *Arch Dis Child* 2000;83:307–12.
146. Gilliland FD, Li YF, Peters JM. Effects of maternal smoking during pregnancy and environmental tobacco smoke on asthma and wheezing in children. *Am J Respir Crit Care Med* 2001;163:429–36.
147. Alexander S, Dodds L, Armson BA. Perinatal outcomes in women with asthma during pregnancy. *Obstet Gynecol* 1998;92:435–40.
148. Kallen B, Rydhstroem H, Aberg A. Asthma during pregnancy—a population based study. *Eur J Epidemiol* 2000;16:167–71.
149. Kramer MS, Coates AL, Michoud MC, et al. Maternal asthma and idiopathic preterm labor. *Am J Epidemiol* 1995;142:1078–88.
150. Perlow JH, Montgomery D, Morgan MA, et al. Severity of asthma and perinatal outcome. *Am J Obstet Gynecol* 1992;167:963–7.
151. Stenius-Aarniala BS, Piirila P, Teramo KA. Asthma and pregnancy: a prospective study of 198 pregnancies. *Thorax* 1988;43:12–18.
152. Doucette JT, Bracken MB. Possible role of asthma in the risk of preterm labor and delivery. *Epidemiology* 1993;4:143–50.

153. Bahna SL, Bjerkedal T. The course and outcome of pregnancy in women with bronchial asthma. *Acta Allergol* 1972;27:397–406.
154. Mabie WC, Barton JR, Sibai BM, et al. Adult respiratory distress syndrome in pregnancy. *Am J Obstet Gynecol* 1992;167:950–7.
155. Liu S, Wen SW, Demissie K, et al. Maternal asthma and pregnancy outcomes: a retrospective cohort study. *Am J Obstet Gynecol* 2001;184:90–6.
156. Demissie K, Breckenridge MB, Rhoads GG. Infant and maternal outcomes in the pregnancies of asthmatic women. *Am J Respir Crit Care Med* 1998;158:1091–5.
157. Schatz M, Zeiger RS, Hoffman CP. Intra-uterine growth is related to gestational pulmonary function in pregnant asthmatic women. *Chest* 1990;98:389–92.
158. Lao TT, Huengsburg M. Labour and delivery in mothers with asthma. *Eur J Obstet Gynecol Reprod Biol* 1990;35:183–90.
159. Corchia C, Bertollini R, Forastiere F, et al. Is maternal asthma a risk factor for low birth weight? Results of an epidemiologic survey. *Eur J Epidemiol* 1995;11:627–31.
160. Gordon M, Niswander KR, Berendes H, et al. Fetal morbidity following potentially anoxigenic obstetric conditions. *Am J Obstet Gynecol* 1970;106:421–9.
161. Greenberger PA, Patterson R. The outcome of pregnancy complicated by severe asthma. *Allergy Proc* 1988;9:539–43.
162. Clifton VL, Giles WB, Smith R, et al. Alterations of placental vascular function in asthmatic pregnancies. *Am J Respir Crit Care Med* 2001;164:546–53.
163. Bjorksten B, Finnstrom O, Wichman K. Intrauterine exposure to the beta-adrenergic receptor-blocking agent metoprolol and allergy. *Int Arch Allergy Immunol* 1988;87:59–62.
164. Nelson H. Pregnancy and allergic diseases. In: Biermaan C, Pearlman, D, eds. *Allergic diseases of infancy, childhood and adolescence*. Philadelphia, PA: W B Saunders, 1980:675–80.
165. Tan KS, Thompson NC. Asthma in pregnancy. *Am J Med* 2000;109:727–33.
166. Kwon HL, Belanger K, Bracken MB. Asthma prevalence during pregnancy in the United States: estimates from national health surveys. *Ann Epidemiol* (in press).
167. Schatz M, Zeiger RS, Hoffman CP, et al. Perinatal outcomes in the pregnancies of asthmatic women: a prospective controlled analysis. *Am J Respir Crit Care Med* 1995;151:1170–4.
168. Jana N, Vasishta K, Saha SC, et al. Effect of bronchial asthma on the course of pregnancy, labour and perinatal outcome. *J Obstet Gynecol* 1995;21:227–32.
169. Chan KN, Elliman A, Bryan E, et al. Respiratory symptoms in children of low birth weight. *Arch Dis Child* 1989;64:1294–304.
170. Chan KN, Noble-Jamieson CM, Elliman A, et al. Lung function in children of low birth weight. *Arch Dis Child* 1989;64:1284–93.
171. O'Shaheen S, Sterne JAC, Montgomery HA. Birth weight, body mass index and asthma in young adults. *Thorax* 1999;54:396–402.
172. Darlow BA, Horwood LJ, Mogridge N. Very low birth weight and asthma by age seven years in a national cohort. *Pediatr Pulmonol* 2000;30:291–6.
173. Nafstad P, Magnus P, Jaakkola JJ. Risk of childhood asthma and allergic rhinitis in relation to pregnancy complications. *J Allergy Clin Immunol* 2000;106:867–73.
174. Pekkanen J, Xu B, Jarvelin MR. Gestational age and occurrence of atopy at age 31: a prospective birth cohort study in Finland. *Clin Exp Allergy* 2001;31:95–102.
175. Rasanen M, Kaprio J, Laitinen T, et al. Perinatal risk factors for asthma in Finnish adolescent twins. *Thorax* 2000;55:25–31.
176. Bjorksten B, Kjellman NIM. Perinatal risk factors influencing the development of allergy. *Clin Exp Allergy* 1990;20:3–8.
177. Xu B, Pekkanen J, Harikainen AL, et al. Caesarean section and risk of asthma and allergy in childhood. *J Allergy Clin Immunol* 2001;107:732–3.
178. Brown MA, Halonen M. Perinatal events in the development of asthma. *Curr Opin Pulm Med* 1999;5:4–9.
179. Xu B, Pekkanen J, Jarvelin MR. Obstetric complications and asthma in childhood. *J Asthma* 2000;37:589–94.
180. Annesi-Maesano I, Moreau D, Strachan D. In utero and perinatal complications preceding asthma. *Allergy* 2001;56:491–7.
181. Gergen PJ, Mullally DI, Evans R III. National survey of prevalence of asthma among children in the United States, 1976–1980. *Pediatrics* 1988;81:1–7.
182. McKeever TM, Lewis SA, Smith C, et al. Siblings, multiple births, and the incidence of allergic disease: a birth cohort study using the West Midlands general practice research database. *Thorax* 2001;56:758–62.
183. Miller JE. Predictors of asthma in young children: does reporting source affect our conclusions? *Am J Epidemiol* 2001;154:245–50.
184. Oddy WH, Holt PG, Sly PD, et al. Association between breast feeding and asthma in 6 year old children: findings of a prospective birth cohort study. *BMJ* 1999;319:815–19.
185. Dell S, To T. Breastfeeding and asthma in young children: findings from a population-based study. *Arch Pediatr Adolesc Med* 2001;155:1261–5.
186. Rusconi F, Galassi C, Corbo GM, et al. Risk factors for early, persistent, and late-onset wheezing in young children. *Am J Respir Crit Care Med* 1999;160:1617–22.
187. Wright AL, Holberg CJ, Taussig LM, et al. Relationship of infant feeding to recurrent wheezing at age 6 years. *Arch Pediatr Adolesc Med* 1995;149:758–63.
188. Sears MR, Greene JM, Willan AR, et al. Long-term relation between breastfeeding and development of atopy and asthma in children and young adults: a longitudinal study. *Lancet* 2002;360:901–7.
189. Wright AL, Holberg CJ, Taussig LM, et al. Factors influencing the relation of infant feeding to asthma and recurrent wheeze in childhood. *Thorax* 2001;56:192–7.
190. Takemura Y, Sakurai Y, Honjo S, et al. Relation between breastfeeding and the prevalence of asthma: the Tokorozawa Childhood Asthma and Pollinosis Study. *Am J Epidemiol* 2001;154:115–19.
191. Burr ML, Limb ES, Maguire MJ, et al. Infant feeding, wheezing, and allergy: a prospective study. *Arch Dis Child* 1993;68:724–8.
192. Wilson AC, Forsyth JS, Greene SA, et al. Relation of infant diet to childhood health: seven year follow up of cohort of children in Dundee infant feeding study. *BMJ* 1998;316:21–5.
193. Delespesse G. Presence of IgE suppressive factors in human colostrum. *Eur J Immunol* 1986;16:1005–8.
194. Lucas A, Brooke OG, Morley R, et al. Early diet of preterm infants and development of allergic or atopic disease: randomized prospective study. *BMJ* 1990;300:837–40.
195. van Asperen PP, Kemp AS, Mellis CM. Immediate food hypersensitivity reaction on the first known exposure to the food. *Arch Dis Child* 1983;58:253–6.
196. Hattevig G, Kjellman B, Johansson SG, et al. Clinical symptoms and IgE responses to common food proteins in atopic and healthy children. *Clin Allergy* 1984;14:551–9.
197. Chandra RK, Puri S, Hamed A. Influence of maternal diet during lactation and use of formula feeds on development of atopic eczema in high risk infants. *BMJ* 1989;299:228–30.

198. Zeiger RS, Heller S. The development and prediction of atopy in high-risk children: follow-up at age 7 years in a prospective randomized study of combined maternal and infant food allergen avoidance. *J Allergy Clin Immunol* 1995;95:1179–90.
199. Arshad SH, Matthews S, Gant C, et al. Effect of allergen avoidance on development of allergic disorders in infancy. *Lancet* 1992;399:1493–7.
200. Hide DW, Matthews S, Matthews L, et al. Effect of allergen avoidance in infancy on allergic manifestations at age two years. *J Allergy Clin Immunol* 1994;93:842–6.
201. Montgomery SM, Wakefield AJ, Morris DL, et al. The initial care of newborn infants and subsequent hay fever. *Allergy* 2000;55:916–22.
202. Martinez FD. Role of viral infections in the inception of asthma and allergies during childhood: could they be protective? *Thorax* 1994;49:1189–91.
203. Strannegard O. Allergy and neonatal care. *Allergy* 2000;55:903–4.
204. Holt PG, Jones CA. The development of the immune system during pregnancy and early life. *Allergy* 2000;55:688–97.
205. Farooqi IS, Hopkin JM. Early childhood infection and atopic disorder. *Thorax* 1998;53:927–32.
206. Wickens K, Pearce N, Crane J, et al. Antibiotic use in early childhood and the development of asthma. *Clin Exp Allergy* 1999;29:766–71.
207. Olivetti JF, Kercsmer CM, Redline S. Pre- and perinatal risk factors for asthma in inner city African-American children. *Am J Epidemiol* 1996;143:570–7.
208. Rona RJ, Gulliford MC, Chinn S. Effects of prematurity and intrauterine growth on respiratory health and lung function in childhood. *BMJ* 1993;306:817–20.
209. Barker DJ, Godfrey KM, Fall C, et al. Relation of birth weight and childhood respiratory infection to adult lung function and death from chronic obstructive airways disease. *BMJ* 1991;303:671–5.
210. Platts-Mills TA, Vervloet D, Thomas WR, et al. Indoor allergens and asthma: report of the Third International Workshop, Cuence, Spain. *J Allergy Clin Immunol* 1997;97:1079–84.
211. Platts-Mills T, Thomas WR, Aalberserc RC, et al. Dust mite allergens and asthma: report of a second international workshop. *J Allergy Clin Immunol* 1992;89:1046–60.
212. Platts-Mills T, Ward GW, Sporik R, et al. Epidemiology of the relationship between exposure to indoor allergens and asthma. *Int Arch Allergy Immunol* 1991;94:339–45.
213. Platts-Mills T, De Weck AL. Dust mite allergens and asthma: a world wide problem. *J Allergy Clin Immunol* 1989;83:416–27.
214. Platts-Mills T, Chapman MD. Dust mites: immunology, allergic disease, and environmental control. *J Allergy Clin Immunol* 1987;80:755–75.
215. Arlian LG, Bernstein IL, Gallagher JS. The prevalence of house dust mites, *Dermatophagoides* spp. and associated environmental conditions in homes in Ohio. *J Allergy Clin Immunol* 1982;69:527–32.
216. Tovey ER, Chapman MD, Platts-Mills T. Mite feces are a major source of house dust allergens. *Nature* 1981;289:592–3.
217. Sicherer SH, Wood RA, Eggleston PA. Determinants of airway responses to cat allergen: comparison of environmental challenge to quantitative nasal and bronchial allergen challenge. *J Allergy Clin Immunol* 1997;99(6 part 1):798–805.
218. Pollart SM, Chapman MD, Fiocco GP, et al. Epidemiology of acute asthma. IgE antibodies to common inhalant allergens as a risk factor for emergency room visits. *J Allergy Clin Immunol* 1989;83:875–82.
219. Sears MR, Herbison GP, Holdaway MD, et al. The relative risks of sensitivity to grass pollen, house dust mite and cat dander in the development of childhood asthma. *Clin Exp Allergy* 1989;19:419–24.
220. Van Metre TE, Marsh DG, Adkinson NFJ. Dose of cat (*Felis domesticus*) allergen1 (Fel d I) that induces asthma. *J Allergy Clin Immunol* 1986;78:62–75.
221. Wood R, Eggleston PA, Lind P, et al. Antigenic analysis of household dust samples. *Am Rev Respir Dis* 1988;137:358–63.
222. Luczynska CM, Li Y, Chapman MD, et al. Airborne concentrations and particle size distribution of allergen derived from domestic cats (*Felis domesticus*). Measurements using cascade impactor, liquid impinger, and a two-site monoclonal antibody assay for Fel d I. *Am Rev Respir Dis* 1990;141:361–7.
223. Ingram JM, Sporik R, Rose G, et al. Quantitative assessment of exposure to dog (Can f 1) and cat (Fel d 1) allergens: relation to sensitization and asthma among children living in Los Alamos, New Mexico. *J Allergy Clin Immunol* 1995;96:449–56.
224. Lind P, Norman PS, Newton M, et al. The prevalence of indoor allergens in the Baltimore area: house dust-mite and animal dander antigens measured by immunochemical techniques. *J Allergy Clin Immunol* 1987;80:541–7.
225. Schou C, Hansen GN, Lintner T, et al. Assay for the major dog allergen Can fl: investigation of house dust samples and commercial dog extracts. *J Allergy Clin Immunol* 1991;88:847–53.
226. Rosenstreich DL, Eggleston P, Kattan M, et al. The role of cockroach allergy and exposure to cockroach allergen in causing morbidity among inner-city children with asthma. *N Engl J Med* 1997;336:1356–63.
227. Swanson MC, Agarwal MK, Reed CE. An immunochemical approach to indoor aeroallergen quantitation with a new volumetric air sampler: studies with mite, roach, cat, mouse, and guinea pig antigens. *J Allergy Clin Immunol* 1985;76:724–9.
228. Kang B, Chang JL. Allergenic impact of inhaled arthropod material. *Clin Rev Allergy* 1985;3:363–75.
229. Call RS, Smith TF, Morris E, et al. Risk factors for asthma in inner city children. *Pediatrics* 1992;121:862–6.
230. Norback D, Bjornsson E, Janson D, et al. Current asthma and biochemical signs of inflammation in relation to building dampness in dwellings. *Int J Tuberc Lung Dis* 1999;3:368–76.
231. Burge YH, Platts-Mills T. Indoor biological aerosols. Research Triangle Park, NC: Office of Health and Environmental Assessment, US Environmental Protection Agency, 1991.
232. Cutten AE, Hasmain SM, Segedin BP, et al. The basidiomycete ganoderma and asthma: collection, quantitation and immunogenicity of the spores. *N Z Med J* 1988;101:361–3.
233. O'Rourke MK, Quackenboss J, Lebowitz MD. Indoor pollen and mold characterization from homes in Tucson, Arizona, USA. In: Indoor air '90: The Fifth International Conference on Indoor Air Quality and Climate, Toronto, Canada (Canada Mortgage and Housing Corp), July 29–August 3, 1990. Vol 2: 9–14.
234. Burge H. Bioaerosols: prevalence and health effects in the indoor environment. *J Allergy Clin Immunol* 1990;86:687–705.
235. Gold DR, Burge HA, Carey V, et al. Predictors of repeated wheeze in the first year of life: the relative roles of cockroach, birth weight, acute lower respiratory illness, and maternal smoking. *Am J Respir Crit Care Med* 1999;160:227–36.
236. Lau S, Illi S, Sommerfeld C, et al. Early exposure to house-dust mite and cat allergens and development of childhood asthma: a cohort study. *Lancet* 2000;356:1392–7.

237. Belanger K, Beckett W, Triche E, et al. Symptoms of wheeze and persistent cough in the first year of life: associations with indoor allergens, air contaminants and maternal history of asthma. *Am J Epidemiol* (in press).
238. Brunekreef B, Dockery DW, Speizer FE, et al. Home dampness and respiratory morbidity in children. *Am Rev Respir Dis* 1989;140:1363–7.
239. Dales RE, Zwanenburg H, Burnett R, et al. Respiratory health effects of home dampness and molds among Canadian children. *Am J Epidemiol* 1991;134:196–203.
240. Park JH, Gold DR, Spiegelman DL, et al. House dust endotoxin and wheeze in the first year of life. *Am J Respir Crit Care Med* 2001;163:322–8.
241. Miller RL, Chew GL, Bell CA, et al. Prenatal exposure, maternal sensitization, and sensitization in utero to indoor allergens in an inner-city cohort. *Am J Respir Crit Care Med* 2001;164:995–1001.
242. Riedler J, Braun-Fahrlander C, Eder W, et al. Exposure to farming in early life and development of asthma and allergy: a cross-sectional survey. *Lancet* 2001;358:1129–33.
243. Braun-Fahrlander C, Riedler J, Herz U, et al. Environmental exposure to endotoxin and its relation to asthma in school-age children. *N Engl J Med* 2002;347:869–77.
244. Gold DR. Environmental tobacco smoke, indoor allergens, and childhood asthma. *Environ Health Perspect* 2000;108:643–51.
245. Cook DG, Strachan DP. Health effects of passive smoking: 3. Parental smoking and prevalence of respiratory symptoms and asthma in school age children. *Thorax* 1997;52:1081–94.
246. Gergen PJ. Environmental tobacco smoke as a risk factor for respiratory disease in children. *Respir Physiol* 2001;128:39–46.
247. US Environmental Protection Agency. Respiratory health effects of passive smoking: lung cancer and other disorders. Washington, DC: Office of Health and Environmental Assessment, Office of Research and Development, 1992. (Publication EPA/600/6-90-006B).
248. London SJ, Gauderman WJ, Avol E, et al. Family history and the risk of early-onset persistent, early-onset transient, and late-onset asthma. *Epidemiology* 2001;12:577–83.
249. Murray AB, Morrison BJ. Passive smoking by asthmatics: its greater effect on boys than on girls and on older than on younger children. *Chest* 1988;94:701–8.
250. O'Connor GT, Weiss ST, Tager IB, et al. The effect of passive smoking on pulmonary function and nonspecific bronchial responsiveness in a population based sample of children and young adults. *Am Rev Respir Dis* 1987;135:800–4.
251. Minor TE, Baker JW, Dick EC. Greater frequency of viral respiratory infections in asthmatic children as compared with their nonasthmatic siblings. *J Pediatr* 1974;85:472–7.
252. Minor TE, Dick EC, Baker JW, et al. Rhinovirus and influenza type A infections as precipitants of asthma. *Am Rev Respir Dis* 1976;113:149–53.
253. Busse W, Gern J, Dick E. The role of respiratory viruses in asthma. *Ciba Found Symp* 1997;206:208–13.
254. Anderson HR, Bland JM, Parel S, et al. The natural history of asthma in childhood. *J Epidemiol Community Health* 1986;40:121–9.
255. Pullan CR, Hey EN. Wheezing, asthma, pulmonary dysfunction 19 years after infection with respiratory syncytial virus in infancy. *BMJ* 1982;248:1665–9.
256. Venn AJ, Yemaneberhan H, Bekele Z, et al. Increased risk of allergy associated with the use of kerosene fuel in the home. *Am J Respir Crit Care Med* 2001;164:1660–4.
257. Jarvis D, Chinn S, Luczynska C, et al. Association of respiratory symptoms and lung function in young adults with use of domestic gas appliance. *Lancet* 1996;347:426–31.
258. Melia RJ, Florey CD, Chinn S, et al. Investigations into the relations between respiratory illness in children, gas cooking and nitrogen dioxide in the UK. *Tokai J Exp Clin Med* 1985;10:375–8.
259. Farrow A, Greenwood R, Preece S, et al. Nitrogen dioxide, the oxides of nitrogen, and infants' health symptoms. ALSPAC Study Team. Avon Longitudinal Study of Pregnancy and Childhood. *Arch Environ Health* 1997;52:189–94.
260. Neas LM, Dockery DW, Ware JH, et al. Association of indoor nitrogen dioxide with respiratory symptoms and pulmonary function in children. *Am J Epidemiol* 1991;134:204–19.
261. Garrett MH, Hooper MA, Hooper BM, et al. Respiratory symptoms in children and indoor exposure to nitrogen dioxide and gas stoves. *Am J Respir Crit Care Med* 1998;158:891–5.
262. Lambert WE, Samet JM, Hunt WC, et al. Nitrogen dioxide and respiratory illness in children. Part II: assessment of exposure to nitrogen dioxide. *Res Rep Health Eff Inst* 1993;June:33–50.
263. Berwick M, Leaderer BP, Stolwijk JA, et al. Lower respiratory symptoms in children exposed to nitrogen dioxide from unvented combustion sources. *Environ Int* 1989;15:369–73.
264. Jones CA, Holloway JA, Warner JO. Does atopic disease start in foetal life? *Allergy* 2000;55:2–10.
265. Bjorksten B. The intrauterine and postnatal environments. *J Allergy Clin Immunol* 1999;104:1119–27.
266. Xu B, Jarvelin MR, Pekkanen J. Prenatal factors and occurrence of rhinitis and eczema among offspring. *Allergy* 1999;54:829–36.
267. Wahn U. What drives the allergic march? *Allergy* 2000;55:591–9.
268. Laitinen T, Daly MJ, Rioux JD, et al. A susceptibility locus for asthma-related traits on chromosome 7 revealed by genome-wide scan in a founder population. *Nat Genet* 2001;28:87–91.
269. Schonberger HJAM, Van Schayck CP. Prevention of asthma in genetically predisposed children in primary care—from clinical efficacy to a feasible intervention programme. *Clin Exp Allergy* 1998;28:1325–31.